

Smoking cessation

This decision aid is to help people decide how to stop smoking, if they decide they want to stop.

The main options are:


- Self-managed quitting or reduction. This means trying to stop smoking, or to cut down, without help from a doctor or other health professional. It could mean using nicotine replacement products (such as chewing gum, sprays, patches, inhalers, and lozenges) that can be bought from a pharmacy, self-help books or websites, or just willpower.
- Health professional-assisted quitting or reduction. This means getting help from a doctor or other health professional to cut down or stop smoking. It could mean taking prescription medicines that can reduce cravings and help people stop smoking, or having counselling, or using nicotine replacement products with support from a health professional.
- Alternative therapy quitting or reduction. This may involve things such as acupuncture, hypnosis, homeopathy, herbal remedies, or reflexology (a kind of massage).
- Continuing to smoke.

People can decide to try more than one option. For example, some people might try nicotine replacement from a pharmacy, along with an alternative therapy.

The decisions that people who smoke need to make include:

- Should I try to stop smoking on my own or get help from a health professional?
- I really want to stop smoking, but I know I'll need help. Should I talk to my doctor about medicines that can help, or try something I can buy over the counter?
- I smoke but I feel okay. Should I carry on smoking?

The right decision for you may change over time. Many people try to stop smoking several times before they succeed. You might need to try several methods before you find one that works for you. This decision aid aims to help you make the right choice for you now, but you may wish to change your mind in future.



What are my options?

Treatment	What is the treatment?
Continuing to smoke	<p>Continuing to smoke means that you haven't tried to stop smoking, or you've tried to stop but haven't managed it yet.</p>
Self-managed quitting or reduction	<p>If you find it hard to stop smoking, it may be because you are addicted to nicotine.</p> <p>Self-managed quitting or reduction means trying to stop smoking or to reduce (cut down) how much you smoke on your own, without help from a doctor or other health professional. This could mean:</p> <ul style="list-style-type: none"> • Using nicotine replacement therapy products that you can buy over the counter from a pharmacy • Using self-help materials, such as books or websites, which can give you information and help about reducing or quitting • Rewarding yourself for smoking less or for stopping altogether • Socialising with other people who are trying to stop smoking • Using electronic cigarettes (so-called e-cigarettes) instead of smoking • Just using your willpower to cut down or stop smoking. <p>You could try any combination of these methods.</p>
Health professional-assisted quitting or reduction	<p>Health professional-assisted quitting or reduction means getting help to stop or cut down smoking from your doctor, or from another health professional.</p> <p>The kind of help you can get from health professionals includes medicines to help you stop smoking, nicotine replacement therapy products (such as gum and patches), with support and advice to help you cut down or stop, counselling to help you stop smoking, or different combinations of these treatments.</p> <p>There are two medicines that you can get with a prescription from your doctor, which may help you stop smoking. They are called bupropion and varenicline. Doctors usually only prescribe bupropion or varenicline as part of a programme to stop smoking. This means that if you want to take these drugs you will need to agree to have counselling.</p>
Alternative therapy quitting or reduction	<p>Alternative treatments are things that your doctor probably won't prescribe, but that you're free to try if you think they might help you. They may not have been studied as thoroughly as other options we look at here. This means there is less evidence about how well they work.</p> <p>These options include acupuncture, hypnosis, hypnotherapy, homeopathy, herbal remedies, and reflexology.</p>

Treatment	What is the effect on stopping smoking?
Continuing to smoke	People who don't take steps to stop smoking are likely to carry on smoking.
Self-managed quitting or reduction	<p>We don't know how well using nicotine replacement therapy products that can be bought over the counter on their own works to help people stop smoking. There hasn't been much good research. We do know that it doesn't work as well as using the same products with the support of a health professional.[1]</p> <p>Between 5 in 100 and 12 in 100 people who try to stop smoking without help from others or from treatments stop smoking for at least six months.[2][3]</p> <p>About 7 in 100 people who use self-help books or CDs stop smoking for at least six months.[4][5][6]</p> <p>There is no good evidence that electronic cigarettes (e-cigarettes) can help people stop smoking.</p>
Health professional-assisted quitting or reduction	<p>Prescription medicines can help people stop smoking. About 22 in 100 people who use bupropion stop smoking for at least six months.[7] About 28 in 100 people who use varenicline stop smoking for at least six months.[8]</p> <p>Nicotine replacement therapy with support from a health professional can help people stop smoking. About 17 in 100 people who use nicotine replacement therapy in this way stop smoking for at least six months.[9]</p> <p>Nicotine replacement therapy with support from a health professional can help people reduce the amount they smoke. About 13 in 100 people who use nicotine replacement in this way manage to cut the number of cigarettes they smoke by half.[10]</p> <p>Counselling can help people stop smoking. Between 11 in 100 and 14 in 100 people who use one-to-one counselling stop smoking for at least six months.[11][12][13] About 19 in 100 people who use group counselling stop smoking for at least six months.[14]</p> <p>Using bupropion as well as nicotine replacement therapy doesn't seem to help any more than just using nicotine replacement therapy.[15]</p> <p>We don't know whether combining nicotine replacement therapy with varenicline or with counselling can help people stop smoking. There hasn't been much research.</p>
Alternative therapy quitting or reduction	There isn't much good evidence that alternative therapies of any kind can help people to reduce how much they smoke or to stop smoking altogether.[16][17][18][19]

Treatment	What is the effect on length of life?
Continuing to smoke	People who continue to smoke are unlikely to live as long as those who stop. On average, people who smoke die 16 years earlier than those who don't smoke.[20]
Self-managed quitting or reduction	People who stop smoking using self-management could live longer. On average, people who give up, using any method, between age 25 and 34 gain an extra 10 years of life. Between age 35 and 44, people gain an extra nine years of life. Between age 45 and 54, you'll gain an extra six years of life. Between age 55 and 64, people gain an extra four years of life.[21]
Health professional-assisted quitting or reduction	People who stop smoking with help from a health professional could live longer. On average, people who give up, using any method, between age 25 and 34 gain an extra 10 years of life. Between age 35 and 44, people gain an extra nine years of life. Between age 45 and 54, you'll gain an extra six years of life. Between age 55 and 64, people gain an extra four years of life.[22]
Alternative therapy quitting or reduction	If alternative treatments help you to stop smoking, you could live longer. On average, people who give up, using any method, between age 25 and 34 gain an extra 10 years of life. Between age 35 and 44, people gain an extra nine years of life. Between age 45 and 54, you'll gain an extra six years of life. Between age 55 and 64, people gain an extra four years of life.[23]

Treatment	What is the effect on how long you can stop smoking for?
Continuing to smoke	People who don't take steps to stop smoking are likely to carry on smoking.
Self-managed quitting or reduction	<p>We don't know how well using nicotine replacement therapy products that can be bought over the counter on their own works to help people stop smoking for at least 12 months. There hasn't been much research. We do know that it doesn't work as well as using nicotine replacement therapy with support from a health professional.[24]</p> <p>Between 5 in 100 and 10 in 100 people who try to stop smoking without help from others or from treatments stop smoking for at least 12 months.[25] [26]</p>
Health professional-assisted quitting or reduction	<p>About 18 in 100 people who use bupropion manage to stop smoking for at least 12 months.[27] About 20 in 100 people who use varenicline manage to stop smoking for at least 12 months.[28]</p> <p>About 16 in 100 people who use nicotine replacement therapy with support from a health professional manage to stop smoking for at least 12 months.[29]</p> <p>We don't know if counselling alone can help people stop smoking for at least 12 months. There hasn't been much research.</p>
Alternative therapy quitting or reduction	We don't know if using alternative treatments can help people stop smoking for at least 12 months. There hasn't been much research.

Treatment	What is the effect on quality of life?
Continuing to smoke	<p>People who continue to smoke are more likely than people who stop smoking to have to spend time in hospital.[30]</p> <p>Continuing to smoke is likely to reduce people's quality of life. People who continue to smoke are more likely to have breathing problems, which will make them less able to do everyday things. These breathing problems may be severe.[31]</p>
Self-managed quitting or reduction	People who stop smoking using self-management are likely to have improved quality of life. When answering quality-of-life questionnaires, people who stop smoking have better scores than people who still smoke. People who stop smoking have fewer coughs and breathing problems than people who still smoke.[32]
Health professional-assisted quitting or reduction	People who stop smoking with help from a health professional are likely to have improved quality of life. When answering quality-of-life questionnaires, people who stop smoking have better scores than people who still smoke. People who stop smoking have fewer coughs and breathing problems than people who still smoke.[33]
Alternative therapy quitting or reduction	<p>People who try acupuncture, reflexology, or hypnosis will need to attend appointments. Acupuncture involves using needles, which some people don't like. Alternative treatments may not be available free on the NHS. People may have to pay for them themselves.</p> <p>People who stop smoking using alternative therapies are likely to have improved quality of life. When answering quality-of-life questionnaires, people who stop smoking have better scores than people who still smoke. People who stop smoking have fewer coughs and breathing problems than people who still smoke.[34]</p>

Treatment	What side effects or complications does the treatment have?
Continuing to smoke	<p>About half of all smokers die of diseases caused by smoking. The main ones are lung cancer, heart disease, and stroke.[35]</p>
Self-managed quitting or reduction	<p>Nicotine patches may cause a mild rash in up to 50 in 100 people.[36]</p> <p>Chewing nicotine gum too fast can cause nausea.[37] Nicotine sprays and inhalers can irritate the nose or throat.[38]</p> <p>Some health authorities have warned that electronic cigarettes (e-cigarettes) may not be safe to use, as they have not been properly tested.[39]</p> <p>People who stop smoking often gain some weight. The average amount that people put on is 3.6 kilograms to 5.9 kilograms (8.0 pounds to 13.0 pounds).[40] This is not always permanent. Some people lose the weight they gain, over time. Weight gain may be reduced while people take nicotine replacement therapy.</p>
Health professional-assisted quitting or reduction	<p>Bupropion can cause difficulty sleeping, nausea, and dry mouth. About 10 in 100 people stop taking bupropion because of side effects.[41]</p> <p>About 1 in 1,000 people who take bupropion have seizures (fits).[42]</p> <p>Varenicline may cause nausea in between 17 in 100 and 44 in 100 people. About 10 in 100 people stop taking varenicline because of nausea.[43]</p> <p>Some people who take varenicline become depressed and think about suicide.[44] We're not yet sure how often this happens. This may be more likely to happen to people who have had mental or psychiatric illnesses in the past, who are given extra support.</p> <p>Nicotine patches may cause a mild rash in up to 50 in 100 people.[45]</p> <p>Chewing nicotine gum too fast can cause nausea.[46] Nicotine sprays and inhalers can irritate the nose or throat.[47]</p> <p>Counselling is not likely to cause side effects.</p> <p>People who stop smoking often gain some weight. The average amount that people put on is 3.6 kilograms to 5.9 kilograms (8.0 pounds to 13.0 pounds).[48] This is not always permanent. Some people lose the weight they gain, over time. Weight gain may be reduced while people take nicotine replacement therapy.</p>
Alternative therapy quitting or reduction	<p>Most alternative treatments are not likely to cause side effects. Herbal remedies such as St John's Wort may interfere with other medicines and cause side effects.</p> <p>People who stop smoking often gain some weight. The average amount that people put on is 3.6 kilograms to 5.9 kilograms (8.0 pounds to 13.0 pounds).[49] This is not always permanent. Some people lose the weight they gain, over time. Weight gain may be reduced while people take nicotine replacement therapy.</p>

What are the pros and cons of each option?

People who want to quit smoking have different experiences about the health problems and views on treatment. Choosing the treatment option that is best for the patient means considering how the consequences of each treatment option will affect their life.

Here are some questions people may want to consider when quitting smoking:

- Is living as long as possible the most important thing to them?
- Do they enjoy smoking too much to want to stop?
- Are they willing to use the treatment most likely to help them stop smoking, even if it causes side effects?
- Do they want a treatment that won't cause a lot of side effects?
- Do they want to avoid putting on weight if they stop smoking?
- Do they want a treatment that means they don't have to spend a lot of time at appointments?
- Do they want a treatment that will ease their withdrawal symptoms the most?
- Do they want a treatment that doesn't involve a lot of pills, medicines, or sprays?
- Do they want a treatment where they get support from other people to help them stop smoking?

How do I get support to help me make a decision that is right for me?

Go to <http://sdm.rightcare.nhs.uk/pda/smoking-cessation/> for more detailed information about treatments for **Smoking Cessation**. People using this type of information say they understand the health problem and treatment choices more clearly, and why one treatment is better for them than another. They also say they can talk more confidently about their reasons for liking or not liking an option with health professionals, friends and family.

You can call our Decision Support service on 0845 450 5851 to speak to a trained Health Coach. The Health Coaches will assist you by:

- Providing you with information
- Helping you to understand your condition
- Recognise what is important to you regarding the outcome of treatment
- Identifying potential solutions
- Encourage you in discussing options with your family
- Transferring skills which will assist you in using the information and resources available to you
- Support you in building confidence in discussing your choice with your doctor
- You may find that this can be achieved in one telephone call with a Health Coach; however, if further calls are required to support you in reaching your decision the Health Coach will schedule these with you.

References

References can be viewed online at <http://sdm.rightcare.nhs.uk/pda/smoking-cessation/references/>